arXiv:1302.2359v2 [math.NT] 23 Feb 2013

BINARY QUADRATIC FORMS AND THE FOURIER COEFFICIENTS OF CERTAIN WEIGHT 1 ETA-QUOTIENTS

ALEXANDER BERKOVICH AND FRANK PATANE

Abstract. We generalize an identity proven by Schoeneberg in 1967. Our identity provides a representation of the most general *eta*-products of weight 1 by binary quadratic forms. We discuss the utility of binary quadratic forms in finding a completion for certain *eta*-quotients. We give a collection of examples which illustrate the use of our general identity in deriving the Fourier coefficients of certain *eta*-quotients.

1. Introduction and Notation

Throughout the paper we assume q is a complex number with $|q| < 1$. We use the standard notations

$$(a;q)_\infty := \prod_{n=0}^{\infty}(1-aq^n) \tag{1.1}$$

and

$$E(q) := (q;q)_\infty. \tag{1.2}$$

Next, we recall the Ramanujan theta function

$$f(a,b) := \sum_{n=-\infty}^{\infty} a^{\frac{n(n+1)}{2}} b^{\frac{n(n-1)}{2}}, \qquad |ab|<1. \tag{1.3}$$

The function $f(a,b)$ satisfies the Jacobi triple product identity [3, Entry 19]

$$f(a,b) = (-a;ab)_\infty(-b;ab)_\infty(ab;ab)_\infty, \tag{1.4}$$

along with

$$f(a,b) = a^{\frac{n(n+1)}{2}} b^{\frac{n(n-1)}{2}} f(a(ab)^n, b(ab)^{-n}), \tag{1.5}$$

where $n \in \mathbb{Z}$ [3, Entry 18]. One may use (1.4) to derive the following special cases:

$$E(q) = f(-q,-q^2) = \sum_{n=-\infty}^{\infty} (-1)^n q^{\frac{n(3n-1)}{2}}, \tag{1.6}$$

$$\phi(q) := f(q,q) = \sum_{n=-\infty}^{\infty} q^{n^2} = \frac{E^5(q^2)}{E^2(q^4)E^2(q)}, \tag{1.7}$$

$$\psi(q) := f(q,q^3) = \sum_{n=-\infty}^{\infty} q^{2n^2-n} = \frac{E^2(q^2)}{E(q)}, \tag{1.8}$$

Date: February 26, 2013.
2010 *Mathematics Subject Classification.* 11B65, 11F03, 11F11, 11F20, 11F27, 11E16, 11E20, 11E25, 14K25.
Key words and phrases. *eta*-quotients, binary quadratic forms, theta series, multiplicative functions, Hecke operators, dissections of q-series.

$$f(q,q^2)=\frac{E^2(q^3)E(q^2)}{E(q^6)E(q)}, \tag{1.9}$$

$$f(q,q^5)=\frac{E(q^{12})E^2(q^2)E(q^3)}{E(q^6)E(q^4)E(q)}. \tag{1.10}$$

Note that (1.6) is the famous Euler pentagonal number theorem. Splitting (1.6) according to the parity of the index of summation, we find

$$E(q)=\sum_{n=-\infty}^{\infty} q^{6n^2-n}-q\sum_{n=-\infty}^{\infty} q^{6n^2+5n}=f(q^5,q^7)-qf(q,q^{11}). \tag{1.11}$$

We will employ the similarly derived relations

$$\phi(q)=\phi(q^9)+2qf(q^3,q^{15}), \tag{1.12}$$

$$\phi(q)=\phi(q^4)+2q\psi(q^8), \tag{1.13}$$

and

$$\psi(q)=f(q^3,q^6)+q\psi(q^9). \tag{1.14}$$

We now recall some notation from elementary number theory. Given an integer n, we use the term $\mathrm{ord}_p(n)$ to represent the unique integer with the properties $p^{\mathrm{ord}_p(n)}\mid n$ and $p^{1+\mathrm{ord}_p(n)}\nmid n$.
For an odd prime p, Legendre's symbol is defined by

$$\left(\frac{n}{p}\right)=\begin{cases}1 & \text{if } n \text{ is a quadratic residue modulo } p \text{ and } p\nmid n,\\ -1 & \text{if } n \text{ is a quadratic nonresidue modulo } p \text{ and } p\nmid n,\\ 0 & \text{if } p\mid n\ .\end{cases}$$

Kronecker's symbol $\left(\frac{n}{m}\right)$ is defined by

$$\left(\frac{n}{m}\right)=\begin{cases}1 & \text{if } m=1\ ,\\ 0 & \text{if } m \text{ is a prime dividing } n,\\ \text{Legendre's symbol} & \text{if } m \text{ is an odd prime,}\end{cases}$$

$$\left(\frac{n}{2}\right)=\begin{cases}0 & \text{if } n \text{ is even },\\ 1 & n\equiv \pm 1 \pmod 8,\\ -1 & n\equiv \pm 3 \pmod 8,\end{cases}$$

and in general $\left(\frac{n}{m}\right)=\prod_{i=1}^{s}\left(\frac{n}{p_i}\right)$, where $m=\prod_{i=1}^{s}p_i$ is a prime factorization of m.

Let $(a,b,c):=ax^2+bxy+cy^2$ be a positive definite binary quadratic form of discriminant $d:=b^2-4ac$. The term "form" will always refer to a positive definite binary quadratic form. The form (a,b,c) has the associated theta series

$$B(a,b,c,q):=\sum_{x,y} q^{ax^2+bxy+cy^2}=\sum_{n\geq 0}(a,b,c,n)q^n. \tag{1.15}$$

Note that $(a,b,c,n)=0$ whenever $n\notin\mathbb{Z}_{\geq 0}$.

We recall some well known facts from the theory of binary quadratic forms. If $\left(\frac{d}{p}\right)=-1$, then no form of discriminant d represents the prime p. If two nonequivalent forms of discriminant d both represent a prime $p\nmid d$, then the forms are inverses [8]. Identifying a form and its inverse, we have at most one form of a given discriminant representing a prime $p\nmid d$. Furthermore, if $(a,b,c,p)>0$ then we must have $(a,b,c,p)=12,8$, or 4 according to whether $d=-3$, $d=-4$, or $d<-4$, respectively [8].

We will need a method to determine which primes p are represpresented by a given form. Besides using basic congruence conditions from genus theory, see [8], we follow [7] and [19] by employing the

Weber class polynomial. For a form (a,b,c) with discriminant d, we let $\tau = \frac{-b+\sqrt{d}}{a}$, and recall that the Weber class polynomial, W_d, is defined as the minimal polynomial of $f(\tau)$, where f is a particular normalized Weber function generating the same class field as Klein's modular function $j(\tau)$. (See [7] for details.)

We will give examples where the factorization pattern of $W_d \pmod{p}$ suffices to determine if (a,b,c) represents p. When the factorization pattern of $W_d \pmod{p}$ does not suffice, we use $\mathrm{rem}(z^p, W_d(z)) \pmod{p}$. Here and throughout the manuscript, $\mathrm{rem}(z^p, W_d(z)) \pmod{p}$ denotes the remainder of z^p upon division by $W_d(z)$ modulo p. If $\left(\frac{d}{p}\right) = 1$, an analysis of $\mathrm{rem}(z^p, W_d(z)) \pmod{p}$ always enables us to determine which form of discriminant d represents p.

An *eta*-quotient is a finite product

$$H(q) := q^j \prod_{n=1}^{N} E^{a_n}(q^{b_n}),$$

where $b_1, \ldots, b_N$ are positive integers with $b_1 < b_2 < \ldots < b_N$; $a_1, \ldots, a_N$ are non-zero integers; and $j = \sum_{n=1}^{N} \frac{b_n a_n}{24} \in \mathbb{Z}$. We call H an *eta*-quotient since $q^{\frac{1}{24}} E(q) = \eta(q)$, where $\eta(q)$ is the Dedekind eta function. The weight of H is defined to be $\sum_{n=1}^{N} \frac{a_n}{2}$. The level of H is defined to be $b_1 \cdot b_N$. H has a Fourier expansion, and we will often write the expansion as

$$H(q) = \sum_{n>0} a(n) q^n.$$

We use $[q^n]H(q)$ to denote the Fourier coefficient $a(n)$, and we call the series $\sum_{n>0} a(n)q^n$ multiplicative when $a(n)$ is multiplicative. It is a continuing area of research to determine the Fourier coefficients of certain *eta*-quotients. (See [1], [5], [6], [12], and [20].)

Let (a,b,c) be a form of discriminant d with $B(a,b,c,q) = \sum_{n\geq 0} h(n)q^n$. In [11] Hecke defines the operator T_p by

$$T_p\left(\sum_{n=0}^{\infty} h(n)q^n\right) = \sum_{n=0}^{\infty} \left[h(pn) + \left(\frac{d}{p}\right) h(n/p)\right] q^n. \tag{1.16}$$

Let S be the set of all linear combinations of theta series associated to the forms of discriminant d. Hecke showed that for $s \in S$, we have $T_p(s) \in S$ [11]. When $T_p(s) = s \cdot \lambda_p$ for some constant λ_p, we call s an eigenform of T_p with eigenvalue λ_p.

If $s = \sum h(n)q^n$ is an eigenform for all Hecke operators, then (1.16) implies

$$\lambda_p h(n) = h(pn) + \left(\frac{d}{p}\right) h(n/p) \tag{1.17}$$

for any positive integer n and any prime p. Taking $n = 1$ yields $\lambda_p \cdot h(1) = h(p)$, and we will always take h appropriately normalized so that $h(1) = 1$. Hence the eigenvalues of all T_p are exactly the evaluations of h at the primes. Equation (1.17) shows h is multiplicative. Moreover, taking $n = p^k$ in (1.17) yields

$$h(p^{k+1}) = h(p)h(p^k) - \left(\frac{d}{p}\right) h(p^{k-1}). \tag{1.18}$$

Even though T_p acts on a given theta series, we may refer to its action on the coefficients of the series. One particular case is $B(a,b,c,q) = \sum_{n\geq 0}(a,b,c,n)q^n$, and we write

$$(a,b,c,n) \xrightarrow{T_p} (a,b,c,np) + \left(\frac{d}{p}\right)(a,b,c,n/p) \tag{1.19}$$

to refer to the action of T_p on the coefficients of $B(a,b,c,q)$.

Given $s, s' \in S$ we say that $s + s'$ is a completion of s if $s + s'$ is a an eigenform for all Hecke operators.

In later sections, we find the Fourier coefficients of certain *eta*-quotients H, by finding a completion $s' + H$ with the additional property that s' and H are congruentially disjoint. The ability to extract the coefficients of H by only employing congruences, contributes to the simplicity of our results. In all cases considered, we are always able to employ congruences to extract the coefficients of our desired *eta*-quotient from its completion.

To find a completion, we use certain multiplicative combinations discussed in [18]. Let $A^n(q)$ be the associated theta series of the binary quadratic form A^n (A composed with itself n times). We employ a result of Sun and Williams [18] to find the following multiplicative linear combinations:

Table 1

$C_5 \cong \langle A \rangle$	$\frac{1}{2}\left(I(q) + \frac{-1+\sqrt{5}}{2}A(q) + \frac{-1-\sqrt{5}}{2}A^2(q)\right)$
	$\frac{1}{2}\left(I(q) + \frac{-1-\sqrt{5}}{2}A(q) + \frac{-1+\sqrt{5}}{2}A^2(q)\right)$
$C_6 \cong \langle A \rangle$	$\frac{1}{2}\left(I(q) - A^2(q) \pm (A(q) - A^3(q))\right)$
$C_8 \cong \langle A \rangle$	$\frac{1}{2}\left(I(q) - A^4(q) \pm \sqrt{2}(A(q) - A^3(q))\right)$
$C_4 \times C_2 \cong \langle A, B \rangle$	$\frac{1}{2}\left(I(q) - A^2(q) \pm (B(q) - BA^2(q))\right)$
$C_4 \times C_4 \cong \langle A, B \rangle$	$\frac{1}{2}(I(q) \pm A(q))^2(I(q) - B^2(q))$

,

where $I(q) := A^0(q)$, and the notation $\langle A_1, A_2, \ldots, A_n \rangle$ is used to denote the group generated by $A_1, \ldots, A_n$ with $|A_1| \geq |A_2| \geq \ldots \geq |A_n|$, where $|A|$ denotes the order of A. Here and throughout, we use C_n to denote the cyclic group of order n. Table 1 lists multiplicative combinations that apply directly to the examples appearing in later sections. We remind the reader that the property of being an eigenform for all Hecke operators is a stronger condition than multiplicativity.

In [15], Schoeneberg proved the identity

$$\frac{B\left(6, 1, \frac{k+1}{24}, q\right) - B\left(6, 5, \frac{k+25}{24}, q\right)}{2} = q^{\frac{k+1}{24}} E(q)E(q^k), \tag{1.20}$$

for $k \equiv -1 \pmod{24}$.
In section 2 we state and prove an extra parameter generalization of (1.20). Our identity provides a representation of the most general *eta*-products of weight 1, by the difference of two binary quadratic forms. Many identities for the weight 1 *eta*-products have previously been considered. (See [2], [4], [9], [14], [15], and [16].) However, we re-emphasize that we consider the most general identities for the weight 1 *eta*-products. Our proof is elementary, in the sense that it makes essential use of dissections of q-series, and does not employ modular form techniques.

Sections 3, 4, and 5, contain a diverse collection of explicit examples in which we derive the Fourier coefficients of certain *eta*-quotients. In section 4, we contrast our multiplicative completion of $q^7 E(q^{12})E(q^{156})$ with that of Gordon and Hughes (see [10]). In section 6, we mention an example that has properties unlike the examples previously seen, and we give concluding remarks.

2. elementary proofs of Theorem 2.1 and Theorem 2.2

Theorem 2.1. *If m, s are positive integers with $24s - m > 0$, then*

$$\frac{B(6m, m, s, q) - B(6m, 5m, s+m, q)}{2} = q^s E(q^m)E(q^{24s-m}).$$

Without loss of generality we assume that m and s are coprime, and hence $(6m, m, s)$ and $(6m, 5m, s+m)$ are primitive forms, but not necessarily reduced. The transformations $(x, y) \to \left(x + \frac{y}{2}, -y\right)$ and $(x, y) \to \left(x + \frac{y}{3}, y\right)$, send $(6m, m, s)$ to $(6m, 5m, s+m)$. Using these transformations we see $(6m, m, s)$ and $(6m, 5m, s+m)$ are equivalent over the p-adic integers for all primes p. Equivalence over the p-adic integers implies $(6m, m, s)$ and $(6m, 5m, s+m)$ share the same genus [7]. Hence

$q^sE(q^m)E(q^{24s-m})$ is a cusp form with simple congruential properties [17, p. 577].
We proceed with the proof of Theorem 2.1.

Proof. For a fixed j we have

$$\sum_{\substack{x\\ y\equiv j \pmod{12}}} q^{6mx^2+mxy+sy^2} = \sum_{\substack{x\\ y\equiv -j \pmod{12}}} q^{6mx^2+mxy+sy^2} = \sum_{x,y} q^{6mx^2+mx(12y+j)+s(12y+j)^2}. \tag{2.1}$$

Letting $x \to -x+y$ and $y \to -y$ on the right of (2.1) yields,

$$\begin{aligned}\sum_{x,y} q^{j^2s-jmx+6mx^2-j(24s-m)y+6(24s-m)y^2} &= q^{sj^2}\left(\sum_x q^{m(6x^2-jx)}\right)\left(\sum_y Q^{(6y^2-jy)}\right)\\ &= q^{sj^2} f\left(q^{m(6-j)},q^{m(6+j)}\right) f\left(Q^{6-j},Q^{6+j}\right),\end{aligned} \tag{2.2}$$

with $Q := q^{24s-m}$.
Employing (2.1) and (2.2) we obtain

$$\begin{aligned}B(6m,m,s,q) =& \sum_{j=0}^{11} q^{sj^2} f\left(q^{m(6-j)},q^{m(6+j)}\right) f\left(Q^{6-j},Q^{6+j}\right)\\ =& f\left(q^{6m},q^{6m}\right) f\left(Q^6,Q^6\right)\\ &+ 2\sum_{j=1}^{5} q^{sj^2} f\left(q^{m(6-j)},q^{m(6+j)}\right) f\left(Q^{6-j},Q^{6+j}\right)\\ &+ q^{36s} f\left(1,q^{12m}\right) f\left(1,Q^{12}\right).\end{aligned} \tag{2.3}$$

We repeat the above process with the form $(6m,5m,s+m)$ to get the analogous seven term expansion for $B(6m,m,s,q)$:

$$\begin{aligned}B(6m,5m,s+m,q) =& \sum_{j=0}^{11} q^{(s+m)j^2} f\left(q^{m(6-5j)},q^{m(6+5j)}\right) f\left(Q^{6-j},Q^{6+j}\right)\\ =& f\left(q^{6m},q^{6m}\right) f\left(Q^6,Q^6\right)\\ &+ 2\sum_{j=1}^{5} q^{(s+m)j^2} f\left(q^{m(6-5j)},q^{m(6+5j)}\right) f\left(Q^{6-j},Q^{6+)}\right)\\ &+ q^{36(s+m)} f\left(q^{-24m},q^{36m}\right) f\left(1,Q^{12}\right).\end{aligned} \tag{2.4}$$

Appropriately applying (1.5) to the right side of (2.4), we write (2.4) as

$$\begin{aligned}B(6m,5m,s+m,q) =& f\left(q^{6m},q^{6m}\right) f\left(Q^6,Q^6\right)\\ &+ 2q^{s+m} f\left(q^{m},q^{11m}\right) f\left(Q^5,Q^7\right)\\ &+ 2\sum_{j=2}^{4} q^{sj^2} f\left(q^{m(6-j)},q^{m(6+j)}\right) f\left(Q^{6-j},Q^{6+j}\right)\\ &+ 2q^{25s-m} f\left(q^{5m},q^{7m}\right) f\left(Q,Q^{11}\right)\\ &+ q^{36s} f\left(1,q^{12m}\right) f\left(1,Q^{12}\right).\end{aligned} \tag{2.5}$$

It is clear that many of the terms in (2.5) also appear in (2.3). Subtracting (2.5) from (2.3) we perform numerous cancellations to obtain

(2.6)

$$\begin{aligned}\frac{B(6m,m,s,q)-B(6m,5m,s+m,q)}{2} =& q^s f\left(q^{5m},q^{7m}\right) f\left(Q^5,Q^7\right) + q^{25s} f\left(q^{m},q^{11m}\right) f\left(Q,Q^{11}\right)\\ &-q^{s+m} f\left(q^{m},q^{11m}\right) f\left(Q^5,Q^7\right) - q^sQ f\left(q^{5m},q^{7m}\right) f\left(Q,Q^{11}\right).\end{aligned}$$

The four terms on the right of (2.6) can be written as the product

$$q^s \cdot \left(f\left(q^{5m}, q^{7m}\right) - q^m f\left(q^m, q^{11m}\right)\right) \cdot \left(f\left(Q^5, Q^7\right) - Qf\left(Q, Q^{11}\right)\right). \tag{2.7}$$

Employing (1.11), we conclude that

$$\frac{B(6m, m, s, q) - B(6m, 5m, s+m, q)}{2} = q^s E(q^m) E(q^{24s-m}). \tag{2.8}$$

□

Once again, the above identity is new although many special cases have been treated previously. In particular, we mention that [2] contains multiple examples with $m > 1$.

Theorem 2.2. *Let d be a discriminant of binary quadratic forms with $16 \mid d$. Take m to be any odd positive divisor of d with $\frac{d}{m}$ coprime to m. Then*

$$\frac{B\left(m, 0, \frac{|d|}{4m}, q\right) - B\left(4m, 4m, \frac{|d|}{16m} + m, q\right)}{2} = q^m \psi(q^{8m}) \phi\left(-q^{\frac{|d|}{16m}}\right).$$

We remark that unlike Theorem 2.1, the forms of Theorem 2.2 are not necessarily in the same genus.

Proof. Let $k := \frac{|d|}{16m}$. We have

$$\begin{aligned} B(m,0,4k,q) - B(4m,4m,k+m,q) &= \sum_{\substack{x \\ y \equiv 0 \pmod 2}} q^{mx^2+ky^2} - \sum_{x,y} q^{m(2x+y)^2+ky^2} \\ &= \sum_{\substack{x \\ y \equiv 0 \pmod 2}} q^{mx^2+ky^2} - \sum_{x \equiv y \pmod 2} q^{mx^2+ky^2} \\ &= \sum_{\substack{x \equiv 1 \pmod 2 \\ y \equiv 0 \pmod 2}} q^{mx^2+ky^2} - \sum_{x \equiv y \equiv 1 \pmod 2} q^{mx^2+ky^2} \\ &= \sum_{x \equiv 1 \pmod 2} q^{mx^2} \left(\sum_{y \equiv 0 \pmod 2} q^{ky^2} - \sum_{y \equiv 1 \pmod 2} q^{ky^2} \right) \\ &= \left(\sum_x q^{m(2x+1)^2} \right) \left(\sum_y (-1)^y q^{ky^2} \right) \\ &= 2q^m \psi(q^{8m}) \phi(-q^k). \end{aligned}$$

□

3. The Fourier Coefficients of $q^2E(q)E(q^{47})$

In this section we determine the Fourier coefficients of $q^2E(q)E(q^{47})$. We have $\mathrm{CL}(-47) \cong C_5$ and the reduced forms are

$\mathrm{CL}(-47) \cong C_5$		$\left(\frac{-47}{p}\right)$
Principal Genus	$(1,1,12), (2,\pm1,6), (3,\pm1,4)$	$+1$

In the above table, p is taken to be coprime to -47 and represented by the given genus.

Taking $m = 1$ and $s = 2$ in Theorem 2.1 yields

$$\sum_{n>0} a(n)q^n := \frac{B(6,1,2,q) - B(6,5,3,q)}{2} = \frac{B(2,1,6,q) - B(3,1,4,q)}{2} = q^2E(q)E(q^{47}). \tag{3.1}$$

One can check $q^2E(q)E(q^{47})$ is not an eigenform for all Hecke operators; hence, we must find a completion for $q^2E(q)E(q^{47})$. Using the first and second row of Table 1, with $A = (2,1,6)$, we find

$$\sum_{n>0} A_1(n)q^n := \frac{B(1,1,12,q)+\frac{\sqrt{5}-1}{2}B(2,1,6,q)+\frac{-\sqrt{5}-1}{2}B(3,1,4,q)}{2}, \tag{3.2}$$

and

$$\sum_{n>0} A_2(n)q^n := \frac{B(1,1,12,q)+\frac{-\sqrt{5}-1}{2}B(2,1,6,q)+\frac{\sqrt{5}-1}{2}B(3,1,4,q)}{2} \tag{3.3}$$

are multiplicative. We show (3.2) and (3.3) are eigenforms for all Hecke operators, and we find their eigenvalues. We then derive a formula for $A_1(n)$ and $A_2(n)$, and exploit

$$[q^n]q^2E(q)E(q^{47}) = \frac{A_1(n)-A_2(n)}{\sqrt{5}}, \tag{3.4}$$

to find the Fourier coefficients of $q^2E(q)E(q^{47})$.

To show (3.2) and (3.3) are eigenforms for T_p, we consider the action of T_p on the forms of discriminant -47. We separate cases according to the value of $\left(\frac{-47}{p}\right)$.

Case 1: $\left(\dfrac{-47}{p}\right) = 1$**.**

Using the notation of (1.19), we obtain

(a,b,c,n)	$(1,1,12,p)>0$	$(3,1,4,p)>0$	$(2,1,6,p)>0$
$(1,1,12,n)\xrightarrow{T_p}$	$2(1,1,12,n)$	$2(3,1,4,n)$	$2(2,1,6,n)$
$(3,1,4,n)\xrightarrow{T_p}$	$2(3,1,4,n)$	$(1,1,12,n)+(2,1,6,n)$	$(3,1,4,n)+(2,1,6,n)$
$(2,1,6,n)\xrightarrow{T_p}$	$2(2,1,6,n)$	$(3,1,4,n)+(2,1,6,n)$	$(1,1,12,n)+(3,1,4,n)$

.

With the aid of the table above, we compute the action of T_p on $A_1(n)$ and $A_2(n)$:

	$(1,1,12,p)>0$	$(3,1,4,p)>0$	$(2,1,6,p)>0$
$A_1(n)\xrightarrow{T_p}$	$2A_1(n)$	$\frac{-\sqrt{5}-1}{2}A_1(n)$	$\frac{\sqrt{5}-1}{2}A_1(n)$
$A_2(n)\xrightarrow{T_p}$	$2A_2(n)$	$\frac{\sqrt{5}-1}{2}A_2(n)$	$\frac{-\sqrt{5}-1}{2}A_2(n)$

.

.

Hence, (3.2) and (3.3) are eigenforms for T_p, with eigenvalues 2, $\frac{\sqrt{5}-1}{2}$, $\frac{-\sqrt{5}-1}{2}$ when $\left(\frac{-47}{p}\right) = 1$.

Case 2: $\left(\dfrac{-47}{p}\right) = -1$**.**

A prime p with $\left(\frac{-47}{p}\right) = -1$ implies $(a,b,c,p)=0$ for any form of discriminant -47. Thus, (3.2) and (3.3) are eigenforms for such T_p with eigenvalue 0.

Case 3: $\left(\dfrac{-47}{p}\right) = 0$**.**

We find that (3.2) and (3.3) are eigenforms for T_{47} with eigenvalue 1.
We have shown (3.2) and (3.3) are eigenforms for all Hecke operators, and have found their corresponding eigenvalues.

We now state criteria to determine when $(a,b,c,p)>0$ for a form of discriminant -47. We are able to distinguish each case by examining the Weber Class polynomial for discriminant -47, $W_{-47}(z) := z^5+2z^4+2z^3+z^2-1$. As mentioned in section 1, we follow the method of [7] to find:

1. $(1,1,12,p)>0$ iff $p=47$ or $\mathrm{rem}(z^p, W_{-47}(z)) \equiv z \pmod{p}$,
2. $(2,1,6,p)>0$ iff $\mathrm{rem}(94z^p, W_{-47}(z)) \equiv$ $(-47+3r_p)z^4+(-5r_p-47)z^3+(-11r_p-47)z^2+(-5r_p-47)z-47-r_p \pmod{p}$,
3. $(3,1,4,p)$ iff $\mathrm{rem}(94z^p, W_{-47}(z)) \equiv$ $(r_p+47)z^4+(47-r_p)z^3+(7r_p+47)z^2+12zr_p-47+5r_p \pmod{p}$,

where r_p is defined by $(r_p)^2 \equiv -47 \pmod{p}$.

Define S_1, S_2, S_3 to be the set of primes $p \neq 47$ represented by $(1,1,12), (2,1,6)$, and $(3,1,4)$, respectively. Let S_4 be the set of primes p with $\left(\frac{-47}{p}\right)=-1$.
We factor a positive integer n as

$$n = 47^{\mathrm{ord}_{47}(n)} \prod_{p_1 \in S_1} p_1^{\mathrm{ord}_{p_1}(n)} \prod_{p_2 \in S_2} p_2^{\mathrm{ord}_{p_2}(n)} \prod_{p_3 \in S_3} p_3^{\mathrm{ord}_{p_3}(n)} \prod_{p_4 \in S_4} p_4^{\mathrm{ord}_{p_4}(n)}. \tag{3.5}$$

Employing (1.18), along with our computed eigenvalues, we obtain

$$A_1(p^\alpha) = \begin{cases} 1 & p=47, \\ 1+\alpha & p \in S_1, \\ U(\alpha) & p \in S_2, \\ V(\alpha) & p \in S_3, \\ \frac{(-1)^\alpha+1}{2} & p \in S_4, \end{cases} \tag{3.6}$$

and

$$A_2(p^\alpha) = \begin{cases} 1 & p=47, \\ 1+\alpha & p \in S_1, \\ V(\alpha) & p \in S_2, \\ U(\alpha) & p \in S_3, \\ \frac{(-1)^\alpha+1}{2} & p \in S_4, \end{cases} \tag{3.7}$$

where

$$U(z) := \frac{\sin(2\pi(z+1)/5)}{\sin(2\pi/5)}, \tag{3.8}$$

and

$$V(z) := \frac{\sin(4\pi(z+1)/5)}{\sin(4\pi/5)}, \tag{3.9}$$

are arithmetic functions of period 5. Using the multiplicativity of $A_1(n)$ and $A_2(n)$, along with (3.6) and (3.7), we have

$$A_1(n) = \prod_{p_1 \in S_1} (1+\mathrm{ord}_{p_1}(n)) \prod_{p_2 \in S_2} U(\mathrm{ord}_{p_2}(n)) \prod_{p_3 \in S_3} V(\mathrm{ord}_{p_3}(n)) \prod_{p_4 \in S_4} \frac{1+(-1)^{\mathrm{ord}_{p_4}(n)}}{2}, \tag{3.10}$$

and

$$A_2(n) = \prod_{p_1 \in S_1} (1+\mathrm{ord}_{p_1}(n)) \prod_{p_2 \in S_2} V(\mathrm{ord}_{p_2}(n)) \prod_{p_3 \in S_3} U(\mathrm{ord}_{p_3}(n)) \prod_{p_4 \in S_4} \frac{1+(-1)^{\mathrm{ord}_{p_4}(n)}}{2}, \tag{3.11}$$

where n is factored as in (3.5). Employing (3.4), (3.10), and (3.11), we obtain

$$a(n) = \tfrac{1}{\sqrt{5}} \prod_{p_1 \in S_1} (1+\mathrm{ord}_{p_1}(n)) \prod_{p_4 \in S_4} \frac{1+(-1)^{\mathrm{ord}_{p_4}(n)}}{2} \cdot P_n, \tag{3.12}$$

where

$$P_n = \prod_{p_2 \in S_2} U(\mathrm{ord}_{p_2}(n)) \prod_{p_3 \in S_3} V(\mathrm{ord}_{p_3}(n)) - \prod_{p_2 \in S_2} V(\mathrm{ord}_{p_2}(n)) \prod_{p_3 \in S_3} U(\mathrm{ord}_{p_3}(n)). \tag{3.13}$$

We explore the functions $U(z)$ and $V(z)$, and find a different representation for both P_n and $a(n)$. Since $U(n)$ and $V(n)$ are periodic, we can tabulate their explicit values

n	$U(n)$	$V(n)$
$n \equiv 0 \pmod 5$	1	1
$n \equiv 1 \pmod 5$	$\frac{\sqrt{5}-1}{2}$	$\frac{-\sqrt{5}-1}{2}$
$n \equiv 2 \pmod 5$	$\frac{-\sqrt{5}+1}{2}$	$\frac{\sqrt{5}+1}{2}$
$n \equiv 3 \pmod 5$	-1	-1
$n \equiv 4 \pmod 5$	0	0

Given a positive integer n, we define r_i to be the number of primes $p_2 \in S_2$ such that $\mathrm{ord}_{p_2}(n) \equiv i \pmod 5$. Similarly, s_i is the number of primes $p_3 \in S_3$ such that $\mathrm{ord}_{p_3}(n) \equiv i \pmod 5$.

We have

$$\prod_{p_2 \in S_2} U(\mathrm{ord}_{p_2}(n)) = \delta_{r_4,0} \cdot (-1)^{r_1+r_3} \cdot \alpha^{r_1+r_2}, \tag{3.14}$$

and

$$\prod_{p_3 \in S_3} V(\mathrm{ord}_{p_3}(n)) = \delta_{s_4,0} \cdot (-1)^{s_1+s_3} \cdot \beta^{s_1+s_2}, \tag{3.15}$$

where $\alpha := \frac{1-\sqrt{5}}{2}$, $\beta := \frac{1+\sqrt{5}}{2}$, and $\delta_{i,j}$ is 1 if $i = j$ and 0 otherwise.

Then

$$\prod_{p_2 \in S_2} U(\mathrm{ord}_{p_2}(n)) \prod_{p_3 \in S_3} V(\mathrm{ord}_{p_3}(n)) = \delta_{r_4+s_4,0} \cdot (-1)^{r_2+r_3+s_1+s_3} \cdot \beta^{s_1+s_2-r_1-r_2}, \tag{3.16}$$

and

$$\prod_{p_2 \in S_2} V(\mathrm{ord}_{p_2}(n)) \prod_{p_3 \in S_3} U(\mathrm{ord}_{p_3}(n)) = \delta_{r_4+s_4,0} \cdot (-1)^{r_2+r_3+s_1+s_3} \cdot \alpha^{s_1+s_2-r_1-r_2}. \tag{3.17}$$

Subtracting (3.17) from (3.16) yields

$$P_n = (-1)^{r_2+r_3+s_1+s_3} \cdot \delta_{r_4+s_4,0} \cdot \left(\beta^{s_1+s_2-r_1-r_2} - \alpha^{s_1+s_2-r_1-r_2}\right). \tag{3.18}$$

We write (3.12) as

(3.19)

$$a(n) = \tfrac{1}{\sqrt{5}} \delta_{r_4+s_4,0} \cdot (-1)^{r_2+r_3+s_1+s_3} \cdot (\beta^{s_1+s_2-r_1-r_2} - \alpha^{s_1+s_2-r_1-r_2}) \cdot \textstyle\prod_{p_4 \in S_4} \dfrac{1+(-1)^{\mathrm{ord}_{p_4}(n)}}{2} \textstyle\prod_{p_1 \in S_1} (1+\mathrm{ord}_{p_1}(n)),$$

which gives an alternate formulation for the Fourier coefficients of $q^2 E(q) E(q^{47})$.

4. Weight 1 eta-products with levels 135, 272, 1872

Applying Theorem 2.1 with $m = 9, s = 1$ yields

$$\frac{B(54,9,1,q) - B(54,45,10,q)}{2} = \frac{B(1,1,34,q) - B(4,3,9,q)}{2} = qE(q^9)E(q^{15}), \tag{4.1}$$

and $m = 3, s = 2$ gives

$$\frac{B(18,3,2,q) - B(18,15,5,q)}{2} = \frac{B(2,1,17,q) - B(5,5,8,q)}{2} = q^2 E(q^3) E(q^{45}). \tag{4.2}$$

We have $\mathrm{CL}(-135) \cong C_6$ and the reduced forms listed according to genus are

$\mathrm{CL}(-135) \cong C_6$		$\left(\frac{p}{5}\right)$	$\left(\frac{p}{3}\right)$
Principal Genus	$(1,1,34)$, $(4,3,9)$, $(4,-3,9)$	$+1$	$+1$
Second Genus	$(5,5,8)$, $(2,1,17)$, $(2,-1,17)$	-1	-1

.

In the above table, p is taken to be coprime to -135 and represented by the given genus.

With the aid of Table 1 with $A = (2,1,17)$ and $\langle A \rangle \cong C_6$, we find

$$\sum_{n>0} a(n)q^n := \frac{B(1,1,34,q) - B(4,3,9,q) + B(2,1,17,q) - B(5,5,8,q)}{2}, \tag{4.3}$$

and

$$\sum_{n>0} b(n)q^n := \frac{B(1,1,34,q) - B(4,3,9,q) - (B(2,1,17,q) - B(5,5,8,q))}{2}, \tag{4.4}$$

are both multiplicative. Note that (4.3) is the sum of (4.1) and (4.2), and that (4.4) is the difference of (4.1) and (4.2). Similar to the previous example, we show (4.3) and (4.4) are eigenforms for all Hecke operators by examining the action of T_p on the forms of discriminant -135. We use the notation of (1.19).

Case 1: $\left(\frac{-135}{p}\right) = 1$**.**

(a,b,c,n)	$(1,1,34,p) > 0$	$(4,3,9,p) > 0$	$(2,1,17,p) > 0$	$(5,5,8,p) > 0$
$(1,1,34,n) \xrightarrow{T_p}$	$2(1,1,34,n)$	$2(4,3,9,n)$	$2(2,1,17,n)$	$2(5,5,8,n)$
$(4,3,9,n) \xrightarrow{T_p}$	$2(4,3,9,n)$	$(1,1,34,n)+(4,3,9,n)$	$(2,1,17,n)+(5,5,8,n)$	$2(2,1,17,n)$
$(2,1,17,n) \xrightarrow{T_p}$	$2(2,1,17,n)$	$(5,5,8,n)+(2,1,17,n)$	$(1,1,34,n)+(4,3,9,n)$	$2(4,3,9,n)$
$(5,5,8,n) \xrightarrow{T_p}$	$2(5,5,8,n)$	$2(2,1,17,n)$	$2(4,3,9,n)$	$2(1,1,34,n)$

.

Using the above table, it is easy to verify (4.3) and (4.4) are eigenforms under T_p with eigenvalues 2 or -2, for a prime p with $\left(\frac{-135}{p}\right) = 1$.

Case 2: $\left(\frac{-135}{p}\right) = -1$**.**

A form (a,b,c) of discriminant -135 has $(a,b,c,p) = 0$ when $\left(\frac{-135}{p}\right) = -1$, and hence (4.3) and (4.4) are eigenforms for such T_p with eigenvalue 0.

Case 3: $\left(\frac{-135}{p}\right) = 0$**.**

(4.3) is an eigenform for T_5 with eigenvalue -1, and (4.4) is an eigenform for T_5 with eigenvalue 1. Both (4.3) and (4.4) are eigenforms for T_3 with eigenvalue 0.

We have shown (4.3) and (4.4) are eigenforms for all Hecke operators, and have found their eigenvalues.

We now state criteria to determine when $(a,b,c,p) > 0$ for a form of discriminant -135. We examine the factorization of the Weber class polynomial $W_{-135}(x) = x^6 - 3x^5 - 3x^4 + 2x^3 + 30x^2 + 33x - 1$ modulo p. Following the method of [7], we find that for a prime p with $\left(\frac{-135}{p}\right) = 1$, we have

1. p is represented by the form $(1,1,34)$ if and only if $W_{-135}(x)$ splits completely modulo p,
2. p is represented by the form $(4,3,9)$ if and only if $W_{-135}(x)$ factors into two irreducible cubic polynomials modulo p,

(3) p is represented by the form $(5, 5, 8)$ if and only if $W_{-135}(x)$ factors into three irreducible quadratic polynomials modulo p,

(4) p is represented by the form $(2, 1, 17)$ if and only if $W_{-135}(x)$ remains irreducible modulo p.

We define S_1, S_2, S_3, S_4 to be the set of primes $p \neq 5$ represented by $(1, 1, 34)$, $(5, 5, 8)$, $(4, 3, 9)$, $(2, 1, 17)$, respectively. We also take S_5 to be the set of primes p with $\left(\frac{-135}{p}\right) = -1$. Employing (1.18) along with the previously computed eigenvalues, we obtain

$$a(p^\alpha) = \begin{cases} 0 & p = 3,\ \alpha > 0, \\ (-1)^\alpha & p = 5, \\ 1 + \alpha & p \in S_1, \\ (-1)^\alpha(1 + \alpha) & p \in S_2, \\ U(\alpha) & p \in S_3, \\ V(\alpha) & p \in S_4, \\ \frac{1+(-1)^\alpha}{2} & p \in S_5, \end{cases} \tag{4.5}$$

and

$$b(p^\alpha) = \begin{cases} 0 & p = 3,\ \alpha > 0, \\ 1 & p = 5, \\ 1 + \alpha & p \in S_1 \cup S_2, \\ U(\alpha) & p \in S_3 \cup S_4, \\ \frac{1+(-1)^\alpha}{2} & p \in S_5, \end{cases} \tag{4.6}$$

where

$$U(z) := \frac{\sin(2\pi(z + 1)/3)}{\sin(2\pi/3)}, \tag{4.7}$$

and

$$V(z) := \frac{\sin(\pi(z + 1)/3)}{\sin(\pi/3)}. \tag{4.8}$$

The functions $U(n)$ and $V(n)$ are periodic and we tabulate their values

n	$U(n)$	$V(n)$
$n \equiv 0 \pmod 6$	1	1
$n \equiv 1 \pmod 6$	-1	1
$n \equiv 2 \pmod 6$	0	0
$n \equiv 3 \pmod 6$	1	-1
$n \equiv 4 \pmod 6$	-1	-1
$n \equiv 5 \pmod 6$	0	0

.

Given a positive integer n we write

$$n = 3^a 5^b \prod_{p_1 \in S_1} p_1^{\operatorname{ord}_{p_1}(n)} \prod_{p_2 \in S_2} p_2^{\operatorname{ord}_{p_2}(n)} \prod_{p_3 \in S_3} p_3^{\operatorname{ord}_{p_3}(n)} \prod_{p_4 \in S_4} p_4^{\operatorname{ord}_{p_4}(n)} \prod_{p_5 \in S_5} p_5^{\operatorname{ord}_{p_5}(n)}, \tag{4.9}$$

where $a = \operatorname{ord}_3(n)$ and $b = \operatorname{ord}_5(n)$. Given the factorization of n in (4.9), we employ (4.5) and (4.6) to find

(4.10)

$$a(n) = (-1)^{b+t} \cdot \delta_{a,0} \cdot \prod_{p \in S_1 \cup S_2} (1 + \operatorname{ord}_p(n)) \prod_{p_3 \in S_3} U(\operatorname{ord}_{p_3}(n)) \prod_{p_4 \in S_4} V(\operatorname{ord}_{p_4}(n)) \prod_{p_5 \in S_5} \frac{1 + (-1)^{\operatorname{ord}_{p_5}(n)}}{2},$$

and

$$b(n) = \delta_{a,0} \cdot \prod_{p \in S_1 \cup S_2} (1 + \mathrm{ord}_p(n)) \prod_{w \in S_3 \cup S_4} U(\mathrm{ord}_w(n)) \prod_{p_5 \in S_5} \frac{1 + (-1)^{\mathrm{ord}_{p_5}(n)}}{2}, \tag{4.11}$$

where t is the number of primes factors of n, counting multiplicity, that are contained in S_2.

Similar to the previous example, we are able to rewrite (4.10) and (4.11) by exploiting the periodicity of $U(z), V(z)$. Define r_i to be the number of primes $p_3 \in S_3$ such that $\mathrm{ord}_{p_3}(n) \equiv i \pmod 3$, and s_i to be the number of primes $p_4 \in S_4$ such that $\mathrm{ord}_{p_4}(n) \equiv i \pmod 6$.

Given the factorization of n in (4.9) we have,

$$\prod_{p_3 \in S_3} U(\mathrm{ord}_{p_3}(n)) = \delta_{r_2,0} \cdot (-1)^{r_1}, \tag{4.12}$$

and

$$\prod_{p_4 \in S_4} V(\mathrm{ord}_{p_4}(n)) = \delta_{s_2+s_5,0} \cdot (-1)^{s_3+s_4}. \tag{4.13}$$

We rewrite (4.10) and (4.11)

$$a(n) = (-1)^{b+t+r_1+s_3+s_4} \cdot \delta_{a+r_2+s_2+s_5,0} \cdot \prod_{p \in S_1 \cup S_2} (1 + \mathrm{ord}_p(n)) \prod_{p_5 \in S_5} \frac{1 + (-1)^{\mathrm{ord}_{p_5}(n)}}{2}, \tag{4.14}$$

and

$$b(n) = (-1)^{r_1+s_1+s_4} \cdot \delta_{a+r_2+s_2+s_5,0} \cdot \prod_{p \in S_1 \cup S_2} (1 + \mathrm{ord}_p(n)) \prod_{p_5 \in S_5} \frac{1 + (-1)^{\mathrm{ord}_{p_5}(n)}}{2}. \tag{4.15}$$

Both (4.10) and (4.14) give the Fourier coefficients of $qE(q^9)E(q^{15}) + q^2E(q^3)E(q^{45})$, and since $qE(q^9)E(q^{15})$ and $q^2E(q^3)E(q^{45})$ have no common terms we can extract the Fourier coefficients of each product by employing congruences. We have

$$[q^n]qE(q^9)E(q^{15}) = \begin{cases} a(n) & n \equiv 1 \pmod 3, \\ 0 & n \equiv 0, 2 \pmod 3, \end{cases}$$

and

$$[q^n]q^2E(q^3)E(q^{45}) = \begin{cases} a(n) & n \equiv 2 \pmod 3, \\ 0 & n \equiv 0, 1 \pmod 3. \end{cases}$$

Comparing (4.15) with (4.14), we get the simple relation

$$b(n) = (-1)^{s_1+s_3+b+t} a(n), \tag{4.16}$$

with t the same as in (4.10).
Using $a(n) + b(n) = [q^n]2qE(q^9)E(q^{15})$ and $a(n) - b(n) = [q^n]2q^2E(q^3)E(q^{45})$ we find

$$[q^n]2qE(q^9)E(q^{15}) = (1 + (-1)^{b+t+s_1+s_3})a(n), \tag{4.17}$$

and

$$[q^n]2q^2E(q^3)E(q^{45}) = (1 - (-1)^{b+t+s_1+s_3})a(n). \tag{4.18}$$

Thus we are able to derive the Fourier coefficients of $qE(q^9)E(q^{15})$ and $q^2E(q^3)E(q^{45})$ by congruences, and by using more than one completion.

4.1. An Eta-Product of Level 272.

Taking $s = 3$ and $m = 4$ in Theorem 2.1 yields

$$\frac{B(24,4,3,q) - B(24,20,7,q)}{2} = \frac{B(3,2,23,q) - B(7,6,11,q)}{2} = q^3 E(q^4)E(q^{68}). \tag{4.19}$$

We have $\mathrm{CL}(-272) \cong C_8$ and the reduced forms listed by genus are

$\mathrm{CL}(-272) \cong C_8$		$\left(\frac{p}{17}\right)$	$\left(\frac{-1}{p}\right)$
Principal Genus	$(1,0,68)$, $(4,0,17)$, $(8,4,9)$, $(8,-4,9)$	$+1$	$+1$
Second Genus	$(3,2,23)$, $(3,-2,23)$, $(7,6,11)$, $(7,-6,11)$	-1	-1

.

In the above table, p is taken to be coprime to -272 and represented by the given genus.

Using Table 1 with $A = (3,2,23)$ and $\langle A\rangle \cong C_8$, we find

$$\sum_{n>0} A_1(n)q^n := \frac{B(1,0,68,q) - B(4,0,17,q) + \sqrt{2}(B(3,2,23,q) - B(7,6,11,q))}{2}, \tag{4.20}$$

and

$$\sum_{n>0} A_2(n)q^n := \frac{B(1,0,68,q) - B(4,0,17,q) - \sqrt{2}(B(3,2,23,q) - B(7,6,11,q))}{2}, \tag{4.21}$$

are both multiplicative. Actually, (4.20) and (4.21) are eigenforms for all Hecke operators as we will soon show.

By employing (1.13) we derive the identity

$$\frac{B(1,0,68,q) - B(4,0,17,q)}{2} = q\phi(q^{68})\psi(q^8) - q^{17}\phi(q^4)\psi(q^{136}). \tag{4.22}$$

We note that the nonzero terms of the Fourier expansion of (4.22) have exponents congruent to 1 (mod 4), while the expansion of $q^3E(q^4)E(q^{68})$ contains only exponents congruent to 3 (mod 4). In other words, congruences can be employed to extract the coefficients of (4.19) from the completions (4.20) and (4.21).

The action of T_p on the forms of discriminant -272 are omitted since the computations are analogous to the examples for discriminant -47 and -135. The action of T_p on $A_1(n)$, $A_2(n)$ is given below for primes p with $\left(\frac{-272}{p}\right) = 1$:

	$(1,0,68,p) > 0$	$(4,0,17,p) > 0$	$(8,4,9,p) > 0$	$(3,2,23,p) > 0$	$(7,6,11,p) > 0$
$A_1(n) \xrightarrow{T_p}$	$2A_1(n)$	$-2A_1(n)$	0	$\sqrt{2}A_1(n)$	$-\sqrt{2}A_1(n)$
$A_2(n) \xrightarrow{T_p}$	$2A_2(n)$	$-2A_2(n)$	0	$-\sqrt{2}A_2(n)$	$\sqrt{2}A_2(n)$

.

If $\left(\frac{-272}{p}\right) = -1$, then $A_1(n)$ and $A_2(n)$ are eigenforms for T_p with eigenvalue 0. $A_1(n)$ and $A_2(n)$ are eigenforms for T_2, and both have eigenvalue 0. $A_1(n)$ and $A_2(n)$ are eigenforms for T_{17}, and both have eigenvalue -1.

Using (1.18) with our computed eigenvalues, we find

$$A_1(p^\alpha) = \begin{cases} 0 & p = 2,\ \alpha > 0, \\ (-1)^\alpha & p = 17, \\ 1 + \alpha & (1,0,68,p) > 0, \\ (-1)^\alpha(1+\alpha) & p \neq 17,\ (4,0,17,p) > 0, \\ U(\alpha) & (3,2,23,p) > 0, \\ V(\alpha) & (7,6,11,p) > 0, \\ (-1)^{\frac{\alpha}{2}}\frac{(-1)^\alpha+1}{2} & (8,4,9,p) > 0, \\ \frac{(-1)^\alpha+1}{2} & \left(\frac{-17}{p}\right) = -1, \end{cases} \tag{4.23}$$

and

$$A_2(p^\alpha) = \begin{cases} 0 & p = 2,\ \alpha > 0, \\ (-1)^\alpha & p = 17, \\ 1+\alpha & (1,0,68,p) > 0, \\ (-1)^\alpha(1+\alpha) & p \neq 17,\ (4,0,17,p) > 0, \\ V(\alpha) & (3,2,23,p) > 0, \\ U(\alpha) & (7,6,11,p) > 0, \\ (-1)^{\frac{\alpha}{2}}\frac{(-1)^\alpha+1}{2} & (8,4,9,p) > 0, \\ \frac{(-1)^\alpha+1}{2} & \left(\frac{-17}{p}\right) = -1, \end{cases} \tag{4.24}$$

with

$$U(z) := \frac{\sin(\pi(z+1)/4)}{\sin(\pi/4)}, \tag{4.25}$$

and

$$V(z) := \frac{\sin(3\pi(z+1)/4)}{\sin(3\pi/4)}. \tag{4.26}$$

The functions $U(n)$ and $V(n)$ are periodic and we tabulate their values in Table 2.

Table 2

n	$U(n)$	$V(n)$
$n \equiv 0 \pmod 8$	1	1
$n \equiv 1 \pmod 8$	$\sqrt{2}$	$-\sqrt{2}$
$n \equiv 2 \pmod 8$	1	1
$n \equiv 3 \pmod 8$	0	0
$n \equiv 4 \pmod 8$	-1	-1
$n \equiv 5 \pmod 8$	$-\sqrt{2}$	$\sqrt{2}$
$n \equiv 6 \pmod 8$	-1	-1
$n \equiv 7 \pmod 8$	0	0

.

We now state criteria to determine when a form of discriminant -272 represents a prime p with $\left(\frac{-272}{p}\right) = 1$. The factorization pattern of $W_{-272}(z) := z^8 - z^4 - 4 \pmod p$ does not distinguish between primes represented by $(3,2,23)$ and $(7,6,11)$. We resort to the method of [7] to find:

1. $(1,0,68,p) > 0$ if and only if $\mathrm{rem}(z^p, W_{-272}(z)) \equiv z \pmod p$,
2. $(4,0,17,p) > 0$ if and only if $p = 17$ or $\mathrm{rem}(z^p, W_{-272}(z)) \equiv -z \pmod p$,
3. $(8,4,9,p) > 0$ if and only if $\mathrm{rem}(68z^p, W_{-272}(z)) \equiv 2r_p z^5 - r_p z \pmod p$,
4. $(3,2,23,p)$ if and only if $\mathrm{rem}(272z^p, W_{-272}(z)) \equiv (r_p - 68)z^7 + (68 - 9r_p)z^3 \pmod p$,
5. $(7,6,11,p)$ if and only if $\mathrm{rem}(272z^p, W_{-272}(z)) \equiv (r_p + 68)z^7 + (-68 - 9r_p)z^3 \pmod p$,

where r_p is defined by $(r_p)^2 \equiv -272 \pmod p$.

Define S_1 to be the set of primes $p \neq 17$ represented by $(1,0,68)$ or $(4,0,17)$. We also define S_2, S_3 to be the set of primes p represented by $(3,2,23)$, $(7,6,11)$, respectively. Lastly, we let S_4 be the set of primes p with $(8,4,9,p) > 0$ or $\left(\frac{-272}{p}\right) = -1$.

Using (4.23) and (4.24) we have

$$A_1(n) = (-1)^{b+t+\frac{s}{2}} \cdot \delta_{a,0} \prod_{p_1 \in S_1} (1+\mathrm{ord}_{p_1}(n)) \prod_{p_2 \in S_2} U(\mathrm{ord}_{p_2}(n)) \prod_{p_3 \in S_3} V(\mathrm{ord}_{p_3}(n)) \prod_{p_4 \in S_4} \frac{(-1)^{\mathrm{ord}_{p_4}(n)}+1}{2}, \tag{4.27}$$

and

$$A_2(n) = (-1)^{b+t+\frac{s}{2}} \cdot \delta_{a,0} \prod_{p_1 \in S_1} (1+\mathrm{ord}_{p_1}(n)) \prod_{p_2 \in S_2} V(\mathrm{ord}_{p_2}(n)) \prod_{p_3 \in S_3} U(\mathrm{ord}_{p_3}(n)) \prod_{p_4 \in S_4} \frac{(-1)^{\mathrm{ord}_{p_4}(n)}+1}{2}, \tag{4.28}$$

where $b = \mathrm{ord}_{17}(n)$, $a = \mathrm{ord}_2(n)$, t is the number of primes factors $p \neq 17$ of n, counting multiplicity, that are represented by $(4, 0, 17)$, and s is the number of primes factors of n, counting multiplicity, that are represented by $(8, 4, 9)$. Note that if s is odd, then $\mathrm{ord}_{p_4}(n)$ is odd for some $p_4 \in S_4$, and thus (4.27) and (4.28) vanish.

As mentioned earlier, we are able to employ congruences to extract the coefficients of $q^3E(q^4)E(q^{68})$ from the completion (4.20). We obtain

$$[q^n]q^3E(q^4)E(q^{68}) = \begin{cases} A_1(n)/\sqrt{2} & n \equiv 3 \pmod 4, \\ 0 & \text{otherwise.} \end{cases}$$

Similar to the examples for discriminant -47 and -135, we are able to rewrite (4.27) and (4.28) by using the properties of $U(z)$ and $V(z)$. Given a positive integer n, we define r_i to be the number of prime factors of n with $p_2 \in S_2$ such that $\mathrm{ord}_{p_2}(n) \equiv i \pmod 8$. Similarly, s_i is the number of prime factors of n with $p_3 \in S_3$ and $\mathrm{ord}_{p_3}(n) \equiv i \pmod 8$.

With the notation of r_i and s_i we have

$$\prod_{p_2 \in S_2} U(\mathrm{ord}_{p_2}(n)) = \delta_{r_3+r_7,0} \cdot (-1)^{r_4+r_5+r_6} \cdot 2^{\frac{r_1+r_5}{2}}, \tag{4.29}$$

and

$$\prod_{p_3 \in S_3} V(\mathrm{ord}_{p_3}(n)) = \delta_{s_3+s_7,0} \cdot (-1)^{s_1+s_4+s_6} \cdot 2^{\frac{s_1+s_5}{2}}, \tag{4.30}$$

where $\delta_{i,j}$ is 1 if $i = j$ and 0 otherwise. Letting $k_i := r_i + s_i$ for $i = 1, 2$, we obtain

$$\prod_{p_2 \in S_2} U(\mathrm{ord}_{p_2}(n)) \prod_{p_3 \in S_3} V(\mathrm{ord}_{p_3}(n)) = \delta_{k_3+k_7,0} \cdot (-1)^{s_1+r_5+k_4+k_6} \cdot 2^{\frac{k_1+k_5}{2}} \tag{4.31}$$

and

$$\prod_{p_2 \in S_2} V(\mathrm{ord}_{p_2}(n)) \prod_{p_3 \in S_3} U(\mathrm{ord}_{p_3}(n)) = \delta_{k_3+k_7,0} \cdot (-1)^{r_1+s_5+k_4+k_6} \cdot 2^{\frac{k_1+k_5}{2}}. \tag{4.32}$$

Using (4.31) and (4.32) we reformulate (4.27) and (4.28) as

$$A_1(n) = (-1)^{b+t+\frac{s}{2}+s_1+r_5+k_4+k_6} \cdot \delta_{a+k_3+k_7,0} \cdot 2^{\frac{k_1+k_5}{2}} \prod_{p_1 \in S_1} (1+\mathrm{ord}_{p_1}(n)) \prod_{p_4 \in S_4} \frac{(-1)^{\mathrm{ord}_{p_4}(n)}+1}{2}, \tag{4.33}$$

and

$$A_2(n) = (-1)^{b+t+\frac{s}{2}+r_1+s_5+k_4+k_6} \cdot \delta_{a+k_3+k_7,0} \cdot 2^{\frac{k_1+k_5}{2}} \prod_{p_1 \in S_1} (1+\mathrm{ord}_{p_1}(n)) \prod_{p_4 \in S_4} \frac{(-1)^{\mathrm{ord}_{p_4}(n)}+1}{2}, \tag{4.34}$$

with a, b, t, s the same as in (4.28).

We now exploit

$$A_1(n) - A_2(n) = [q^n]2\sqrt{2}q^3E(q^4)E(q^{68}), \tag{4.35}$$

along with (4.27), and (4.28), to write a formula for $[q^n]q^3E(q^4)E(q^{68})$ without reference to congruences.

Comparing (4.33) and (4.34), we see the relation,

$$A_1(n) = (-1)^{k_1+k_5}A_2(n). \tag{4.36}$$

Combining (4.36) and (4.35), we obtain

$$[q^n]2\sqrt{2}q^3E(q^4)E(q^{68}) = A_1(n)(1-(-1)^{k_1+k_5}), \tag{4.37}$$

and thus the Fourier coefficients of $2\sqrt{2}q^2E(q^4)E(q^{68})$ are given by

(4.38)

$$(1-(-1)^{k_1+k_5})(-1)^{b+t+\frac{s}{2}+s_1+r_5+k_4+k_6}\cdot\delta_{a+k_3+k_7,0}\cdot 2^{\frac{k_1+k_5}{2}}\prod_{p_1\in S_1}(1+\mathrm{ord}_{p_1}(n))\prod_{p_4\in S_4}\frac{(-1)^{\mathrm{ord}_{p_4}(n)}+1}{2}.$$

4.2. **An Eta-Product of Level 1872.** Applying Theorem 2.1 with $m = 12$ and $s = 7$, yields

$$\frac{B(72,12,7,q)-B(72,60,19,q)}{2} = \frac{B(7,2,67,q)-B(19,16,28,q)}{2} = q^7E(Q)E(Q^{13}), \tag{4.39}$$

where we define $Q := q^{12}$.

We have $\mathrm{CL}(-1872) \cong C_4 \times C_4$ and

$\mathrm{CL}(-1872) \cong C_4 \times C_4$		$\left(\frac{p}{3}\right)$	$\left(\frac{p}{13}\right)$	$\left(\frac{-1}{p}\right)$
Principal Genus	$(1,0,468)$, $(4,0,117)$,$(9,0,52)$,$(13,0,36)$	$+1$	$+1$	$+1$
Second Genus	$(7,2,67)$, $(7,-2,67)$, $(19,16,28)$, $(19,-16,28)$	$+1$	-1	-1
Third Genus	$(8,4,59)$, $(8,-4,59)$, $(11,8,44)$, $(11,-8,44)$	-1	-1	-1
Fourth Genus	$(9,6,53)$, $(9,-6,53)$, $(17,10,29)$, $(17,-10,29)$	-1	$+1$	$+1$

In the above table, p is taken to be coprime to -1872 and represented by the given genus.

Using the bottom row of Table 1 with $A = (7,2,67), B = (11,8,44)$ we find

(4.40)

$$\sum_{n>0} A_1(n)q^n := \frac{B(1,0,468,q)+B(13,0,36,q)-B(4,0,117,q)-B(9,0,52,q)}{2} + 2q^7E(Q)E(Q^{13}),$$

and

(4.41)

$$\sum_{n>0} A_2(n)q^n := \frac{B(1,0,468,q)+B(13,0,36,q)-B(4,0,117,q)-B(9,0,52,q)}{2} - 2q^7E(Q)E(Q^{13}).$$

are multiplicative.

Appropriately applying (1.12), (1.13), and (1.14) to

$$\frac{B(1,0,468,q)+B(13,0,36,q)-B(4,0,117,q)-B(9,0,52,q)}{2}, \tag{4.42}$$

we see wonderful cancellation that transforms (4.42) into

$$q\left[\phi(Q^{39})f(Q^2,Q^4)+Q\phi(Q^3)f(Q^{26},Q^{52})-2Q^5\psi(Q^6)f(Q^{13},Q^{65})-2Q^{10}\psi(Q^{78})f(Q,Q^5)\right]. \tag{4.43}$$

Since $Q := q^{12}$, the Fourier expansion of (4.43) only contains terms with exponents congruent to $1 \pmod{12}$. Similarly, the Fourier expansion of $q^7E(Q)E(Q^{13})$ only contains terms with exponents congruent to $7 \pmod{12}$. Hence congruences can be employed to extract coefficients of (4.39) from the completion (4.40).

In [10], Gordon and Hughes consider the product $q^7E(Q)E(Q^{13})$, and introduce the function

(4.44)

$$h(q) = q\left[\phi(-Q^{39})f(Q^2,Q^4)+Q\phi(-Q^3)f(Q^{26},Q^{52})-Q^5\psi(Q^6)f(Q^{13},Q^{65})-Q^{10}\psi(Q^{78})f(Q,Q^5)\right].$$

One can verify $h(q) + 2q^7E(Q)E(Q^{13})$ is an eigenform for some, but not all Hecke operators. Indeed, $h(q) + 2q^7E(Q)E(Q^{13})$ is not an eigenform for T_7, T_{11}, T_{17}, among others. We note the striking similarlity between (4.43) and $h(q)$.

The action of T_p on the forms of discriminant -1872 are omitted since the computations are analogous to previous examples. The action of T_p on $A_1(n)$ and $A_2(n)$ is given below for primes p with $\left(\frac{p}{3}\right) = \left(\frac{-13}{p}\right) = 1$. Note the property $\left(\frac{p}{3}\right) = \left(\frac{-13}{p}\right) = 1$ is equivalent to the prime $p \neq 13$ being represented by the principal genus or second genus, acoording to the table of genera at the beginning of this example.

	$(1,0,468,p)>0$	$(4,0,117,p)>0$	$(13,0,36,p)>0$	$(9,0,52,p)>0$	$(7,2,67,p)>0$	$(19,16,28,p)>0$
$A_1(n) \xrightarrow{T_p}$	$2A_1(n)$	$-2A_1(n)$	$2A_1(n)$	$-2A_1(n)$	$2A_1(n)$	$-2A_1(n)$
$A_2(n) \xrightarrow{T_p}$	$2A_2(n)$	$-2A_2(n)$	$2A_2(n)$	$-2A_2(n)$	$-2A_2(n)$	$2A_2(n)$

.

.

If $p \neq 13$ does not have $\left(\frac{p}{3}\right) = \left(\frac{-13}{p}\right) = 1$, then $A_1(n)$ and $A_2(n)$ are eigenforms for T_p with eigenvalue 0. $A_1(n)$ and $A_2(n)$ are eigenforms for T_{13} with eigenvalue 1. Thus $A_1(n)$ and $A_2(n)$ are eigenforms for all Hecke operators. Employing (1.18) we obtain

$$A_1(p^\alpha) = \begin{cases} 0 & p = 2,3,\ \alpha > 0, \\ 1 & p = 13, \\ 1+\alpha & p \neq 13, (1,0,468,p)+(13,0,36,p)+(7,2,67,p) > 0, \\ (-1)^\alpha(1+\alpha) & (4,0,117,p)+(9,0,52,p)+(19,16,28,p) > 0, \\ (-1)^{\frac{\alpha}{2}}\frac{(-1)^\alpha+1}{2} & -\left(\frac{p}{3}\right) = \left(\frac{-13}{p}\right) = 1, \\ \frac{(-1)^\alpha+1}{2} & \left(\frac{-1872}{p}\right) = -1, \end{cases} \tag{4.45}$$

and

$$A_2(p^\alpha) = \begin{cases} 0 & p = 2,3,\ \alpha > 0, \\ 1 & p = 13, \\ 1+\alpha & p \neq 13, (1,0,468,p)+(13,0,36,p)+(19,16,28,p) > 0, \\ (-1)^\alpha(1+\alpha) & (4,0,117,p)+(9,0,52,p)+(7,2,67,p) > 0, \\ (-1)^{\frac{\alpha}{2}}\frac{(-1)^\alpha+1}{2} & -\left(\frac{p}{3}\right) = \left(\frac{-13}{p}\right) = 1, \\ \frac{(-1)^\alpha+1}{2} & \left(\frac{-1872}{p}\right) = -1. \end{cases} \tag{4.46}$$

We notice $|A_1(p^\alpha)| = |A_2(p^\alpha)|$, and to determine the sign we need a way to distinguish between representations by forms of discriminant -1872. Since $|\mathrm{CL}(-1872)| = 16$, the associated Weber class polynomial, $W_{-1872}(x)$, is a degree 16 polynomial. Explicitly,

$$\begin{aligned} W_{-1872}(x) :=& x^{16} - 8x^{15} + 24x^{14} - 34x^{13} + x^{12} + 246x^{11} - 1094x^{10} + 2574x^9 - 4200x^8 + 5608x^7 \\ & - 5144x^6 + 858x^5 + 4189x^4 - 5166x^3 + 2814x^2 - 750x + 69. \end{aligned}$$

Following [7], one can use the remainder criteria that we have seen in previous examples to determine which primes are represented by a form of discriminant -1872. The coefficients in this criteria become rather large and so we omit the computations. We remark that the procedure is completely analogous to that of section 3.
Let S_1 be the set of primes p with $\left(\frac{p}{3}\right) = \left(\frac{-13}{p}\right) = 1$.

Employing (4.45) and (4.46) we obtain

$$A_1(n) = (-1)^{t_1+\frac{s}{2}} \cdot \delta_{\mathrm{ord}_2(n)+\mathrm{ord}_3(n),0} \prod_{p_1 \in S_1} (1+\mathrm{ord}_{p_1}(n)) \prod_{p_2 \notin S_1 \cup \{2,3,13\}} \frac{(-1)^{\mathrm{ord}_{p_2}(n)}+1}{2}, \tag{4.47}$$

and

$$A_2(n) = (-1)^{t_2+\frac{s}{2}} \cdot \delta_{\mathrm{ord}_2(n)+\mathrm{ord}_3(n),0} \prod_{p_1 \in S_1} (1+\mathrm{ord}_{p_1}(n)) \prod_{p_2 \notin S_1 \cup \{2,3,13\}} \frac{(-1)^{\mathrm{ord}_{p_2}(n)}+1}{2}, \tag{4.48}$$

where n has t_1 prime factors with $(4,0,117,p)+(9,0,52,p)+(19,16,28,p)>0$, t_2 prime factors with $(4,0,117,p)+(9,0,52,p)+(7,2,67,p)>0$, s prime factors with $-\left(\frac{p}{3}\right)=\left(\frac{-13}{p}\right)=1$, and the right most product is taken over primes $p_2\notin S_1\cup\{2,3,13\}$. All prime factors are counted with multiplicity, and similar to the previous example, s odd implies (4.47) and (4.48) vanish.

As mentioned earlier, we can write the Fourier coefficients of $q^7E(Q)E(Q^{13})$ by employing congruences. We have

$$[q^n]q^7E(Q)E(Q^{13})=\begin{cases}\frac{A_1(n)}{2} & n\equiv 7 \pmod{12},\\ 0 & \text{otherwise.}\end{cases}$$

Comparing (4.47) and (4.48), we see

$$A_1(n)=(-1)^{t_1+t_2}A_2(n). \tag{4.49}$$

Subtracting (4.41) from (4.40) gives

$$A_1(n)-A_2(n)=[q^n]4q^7E(Q)E(Q^{13}). \tag{4.50}$$

Combining (4.49) and (4.50) we have

$$[q^n]4q^7E(Q)E(Q^{13})=A_1(n)(1-(-1)^{t_1+t_2}), \tag{4.51}$$

and we can write the Fourier coefficients of $4q^7E(Q)E(Q^{13})$ as

$$(1-(-1)^{t_1+t_2})(-1)^{t_1+\frac{s}{2}}\cdot\delta_{\mathrm{ord}_2(n)+\mathrm{ord}_3(n),0}\prod_{p_1\in S_1}(1+\mathrm{ord}_{p_1}(n))\prod_{p_2\notin S_1\cup\{2,3,13\}}\frac{(-1)^{\mathrm{ord}_{p_2}(n)}+1}{2}. \tag{4.52}$$

5. Certain eta-quotients from the list of Yves Martin

In the following examples we proceed similarly to the examples of section 4, except we refer to [13] to prove the quotient in question is an eigenform for all Hecke operators. We then compute the eigenvalues and the Fourier coefficients of the related quotient.

5.1. An Eta-Quotient of Level 256.

Taking $m=1$ and $d=-256$ in Theorem 2.2, we obtain

$$\sum_{n>0}a(n)q^n:=\frac{B(1,0,64,q)-B(4,4,17,q)}{2}=q\psi(q^8)\phi(-q^{16})=q\frac{E^4(q^{16})}{E(q^8)E(q^{32})}. \tag{5.1}$$

We find $\mathrm{CL}(-256)\cong C_4$ and we list the reduced forms according to their genus

$\mathrm{CL}(-256)\cong C_4$		$\left(\frac{-1}{p}\right)$	$\left(\frac{2}{p}\right)$
Principal Genus	$(1,0,64)$, $(4,4,17)$	$+1$	$+1$
Second Genus	$(5,2,13)$, $(5,-2,13)$	$+1$	-1

In the above table, p is taken to be coprime to -256 and represented by the given genus.

In [13], Y. Martin proves (5.1) is an eigenform for all Hecke operators, but does not give the corresponding eigenvalues $a(p)$. The representation of $q\psi(q^8)\phi(-q^{16})$ in (5.1) allows us to calculate the eigenvalues $a(p)$.

Using $a(p)=\frac{(1,0,64,p)-(4,4,17,p)}{2}$, along with the basic theory of binary quadratic forms, see section 1, we have

$$a(p)=\begin{cases}0 & p=2,\\ 2 & (1,0,64,p)>0,\\ -2 & (4,4,17,p)>0,\\ 0 & p\equiv 5 \pmod 8,\\ 0 & p\equiv 3 \pmod 4).\end{cases} \tag{5.2}$$

Similar to the previous examples, we may use the Weber class polynomial $W_{-256}(x)=x^4-2$ to determine exactly when $(1,0,64,p)>0$ or $(4,4,17,p)>0$. However, we choose to employ an equivalent criteria which does not refer to $W_{-256}(x)$. Following [7], we find that for a prime p we have

(1) $(1,0,64,p)>0$ if and only if $p\equiv 1 \pmod 8$ and $2^{\frac{p-1}{4}}\equiv 1 \pmod p$,

(2) $(4,4,17,p)>0$ if and only if $p\equiv 1 \pmod 8$ and $2^{\frac{p-1}{4}}\equiv -1 \pmod p$.

Employing (1.18) along with (5.2), we find

$$a(p^\alpha)=\begin{cases} 0 & p=2,\ \alpha>0,\\ 1+\alpha & p\equiv 1 \pmod 8 \text{ and } 2^{\frac{p-1}{4}}\equiv 1 \pmod p,\\ (-1)^\alpha(1+\alpha) & p\equiv 1 \pmod 8 \text{ and } 2^{\frac{p-1}{4}}\equiv -1 \pmod p,\\ (-1)^{\frac{\alpha}{2}}\frac{(-1)^\alpha+1}{2} & p\equiv 5 \pmod 8,\\ \frac{(-1)^\alpha+1}{2} & p\equiv 3 \pmod 4.\end{cases} \tag{5.3}$$

To calculate $a(n)$ we partition the odd prime divisors of n according to their residue modulo 8. We write

$$n=2^a\prod_{p_1\equiv 1 \pmod 8} p_1^{\mathrm{ord}_{p_1}(n)}\prod_{p_2\equiv 3,5,7 \pmod 8} p_2^{\mathrm{ord}_{p_2}(n)} \tag{5.4}$$

where $a=\mathrm{ord}_2(n)$.
Using (5.3) and the factorization in (5.4), we derive

$$a(n)=(-1)^{t+\frac{s}{2}}\cdot\delta_{a,0}\prod_{p_1\equiv 1 \pmod 8}(1+\mathrm{ord}_{p_1}(n))\prod_{p_2\equiv 3,5,7 \pmod 8}\frac{(-1)^{\mathrm{ord}_{p_2}(n)}+1}{2}, \tag{5.5}$$

where t is the number of prime factors of n, counting multiplicity, that are congruent to $1 \pmod 8$ and have $2^{\frac{p-1}{4}}\equiv -1 \pmod p$, and s is the number of prime factors of n, counting multiplicity, that are congruent to $5 \pmod 8$. Similar to previous examples, if s is odd then $\mathrm{ord}_{p_2}(n)$ is odd for some $p_2\equiv 5 \pmod 8$, and thus $a(n)$ vanishes.
We remark that formula (5.5) was first stated in [1] without proof.

5.2. **An Eta-Product of Level 80.**

Applying Theorem 2.1 with $m=4$ and $s=1$ yields

$$\sum_{n>0}a(n)q^n:=\frac{B(24,4,1,q)-B(24,20,5,q)}{2}=\frac{B(1,0,20,q)-B(4,0,5,q)}{2}=qE(q^4)E(q^{20}). \tag{5.6}$$

We have $\mathrm{CL}(-80)\cong C_4$, and we explicitly list the reduced forms according to their genus.

$\mathrm{CL}(-80)\cong C_4$		$\left(\frac{p}{5}\right)$	$\left(\frac{-1}{p}\right)$
Principal Genus	$(1,0,20)$, $(4,0,5)$	$+1$	$+1$
Second Genus	$(3,2,7)$, $(3,-2,7)$	-1	-1

In the above table, p is taken to be coprime to -80 and represented by the given genus.

Similar to the previous example, $qE(q^4)E(q^{20})$ is an eigenform for all Hecke operators by appealing to [13]. The representation of $qE(q^4)E(q^{20})$ in (5.6) allows us to calculate the corresponding eigenvalues $a(p)$. Using $a(p)=\frac{(1,0,20,p)-(4,0,5,p)}{2}$, we have

$$a(p)=\begin{cases} 0 & p=2,\\ -1 & p=5,\\ 2 & (1,0,20,p)>0,\\ -2 & p\neq 5, (4,0,5,p)>0,\\ 0 & p\equiv 3,7,11,13,17,19 \pmod{20}.\end{cases} \tag{5.7}$$

To determine whether a given prime $p\equiv 1,9 \pmod{20}$ is represented by $(1,0,20)$ or $(4,0,5)$, one may examine the factorization of the Weber class polynomial $W_{-80}(x)=x^4-5$ modulo p. Like the previous example, we give an equivalent criteria that avoids reference to $W_{-80}(x)$. Following [7], we find that for a prime p we have

(1) p is represented by the form $(1,0,20)$ if and only if
$$(p \equiv 1 \pmod{20} \text{ and } 5^{\frac{p-1}{4}} \equiv 1 \pmod{p}) \text{ or } (p \equiv 9 \pmod{20} \text{ and } 5^{\frac{p-1}{4}} \equiv -1 \pmod{p}),$$

(2) $p \neq 5$ is represented by the form $(4,0,5)$ if and only if
$$(p \equiv 1 \pmod{20} \text{ and } 5^{\frac{p-1}{4}} \equiv -1 \pmod{p}) \text{ or } (p \equiv 9 \pmod{20} \text{ and } 5^{\frac{p-1}{4}} \equiv 1 \pmod{p}).$$

Define S_1, S_2 to be the set of primes $p \neq 5$ represented by $(1,0,20)$, $(4,0,5)$, respectively. Using (1.18) along with (5.7), we find

$$a(p^\alpha) = \begin{cases} 0 & p = 2,\ \alpha > 0, \\ (-1)^\alpha & p = 5, \\ 1+\alpha & p \in S_1, \\ (-1)^\alpha(1+\alpha) & p \in S_2, \\ (-1)^{\frac{\alpha}{2}} \frac{1+(-1)^\alpha}{2} & p \equiv 3, 7 \pmod{20}, \\ \frac{1+(-1)^\alpha}{2} & p \equiv 11, 13, 17, 19 \pmod{20}. \end{cases} \tag{5.8}$$

Employing (5.8), we derive

$$a(n) = (-1)^{t+\frac{s}{2}+\mathrm{ord}_5(n)} \cdot \delta_{\mathrm{ord}_2(n),0} \prod_{p \in S_1 \cup S_2} (1 + \mathrm{ord}_p(n)) \prod_{q \equiv 3,7,11,13,17,19 \pmod{20}} \frac{(-1)^{\mathrm{ord}_q(n)} + 1}{2},$$

where s is the number of prime factors of n, counting multiplicity, congruent to 3,7 (mod 20), and t is the number of prime factors of n, counting multiplicity, contained in S_2. If s is odd, then $\mathrm{ord}_q(n)$ is odd for some $q \equiv 3, 7 \pmod{20}$, and thus $a(n)$ vanishes.

5.3. **Eta-Quotients of Level 144 and 576.** Applying Theorem 2.1 with $s = 1$ and $m = 12$, and reducing the quadratic forms, we obtain

$$\sum_{n>0} a(n)q^n := \frac{B(1,0,36,q) - B(4,0,9,q)}{2} = qE^2(q^{12}). \tag{5.9}$$

We have $\mathrm{CL}(-144) \cong C_4$, and we list the reduced forms according to their genus

$\mathrm{CL}(-144) \cong C_4$		$\left(\frac{p}{3}\right)$	$\left(\frac{-1}{p}\right)$
Principal Genus	$(1,0,36)$, $(4,0,9)$	$+1$	$+1$
Second Genus	$(5,4,8)$, $(5,-4,8)$	-1	$+1$

.

In the above table, p is taken to be coprime to -144 and represented by the given genus.

In [13], Y. Martin proves (5.9) is an eigenform for all Hecke operators.
Using $a(p) = \frac{(1,0,36,p)-(4,0,9,p)}{2}$, we have

$$a(p) = \begin{cases} 0 & p = 2, 3, \\ 2 & (1,0,36,p) > 0, \\ -2 & (4,0,9,p) > 0, \\ 0 & p \equiv 5, 7, 11 \pmod{12}. \end{cases} \tag{5.10}$$

We may use the Weber class polynomial $W_{-144}(x) := x^4 - 2x^3 + 4x - 2$ to determine exactly when $(1,0,36,p) > 0$ or $(4,0,9,p) > 0$. Once again, we employ an equivalent criteria which does not refer to $W_{-144}(x)$. Following [7], we find that for a prime p we have

(1) $(1,0,36,p) > 0$ if and only if $p \equiv 1 \pmod{12}$ and $12^{\frac{p-1}{4}} \equiv 1 \pmod{p}$,

(2) $(4,0,9,p) > 0$ if and only if $p \equiv 1 \pmod{12}$ and $12^{\frac{p-1}{4}} \equiv -1 \pmod{p}$.

Employing (1.18) along with (5.10) we find

$$a(p^\alpha) = \begin{cases} 0 & p = 2, 3\ \alpha > 0, \\ 1+\alpha & p \equiv 1 \pmod{12} \text{ and } 12^{\frac{p-1}{4}} \equiv 1 \pmod{p}, \\ (-1)^\alpha (1+\alpha) & p \equiv 1 \pmod{12} \text{ and } 12^{\frac{p-1}{4}} \equiv -1 \pmod{p}, \\ (-1)^{\frac{\alpha}{2}} \frac{(-1)^\alpha + 1}{2} & p \equiv 5 \pmod{12}, \\ \frac{(-1)^\alpha+1}{2} & p \equiv 7, 11 \pmod{12}. \end{cases} \tag{5.11}$$

To calculate $a(n)$ we partition the prime divisors $p \neq 2, 3$ of n according to their residue modulo 12. We write

$$n = 2^a 3^b \prod_{p_1 \equiv 1 \pmod{12}} p_1^{\operatorname{ord}_{p_1}(n)} \prod_{p_2 \equiv 5,7,11 \pmod{12}} p_2^{\operatorname{ord}_{p_2}(n)}, \tag{5.12}$$

where $a = \operatorname{ord}_2(n)$ and $b = \operatorname{ord}_3(n)$.

Using (5.11) and the factorization in (5.12), we derive

$$a(n) = (-1)^{t+\frac{s}{2}} \cdot \delta_{a+b,0} \prod_{p_1 \equiv 1 \pmod{12}} (1 + \operatorname{ord}_{p_1}(n)) \prod_{p_2 \equiv 5,7,11 \pmod{12}} \frac{(-1)^{\operatorname{ord}_{p_2}(n)} + 1}{2}, \tag{5.13}$$

where t is the number of prime factors of n, counting multiplicity, that are congruent to $1 \pmod{12}$ and have $12^{\frac{p-1}{4}} \equiv -1 \pmod{p}$, and s is the number of prime factors of n, counting multiplicity, that are congruent to $5 \pmod{12}$. We note that s odd implies $\operatorname{ord}_{p_2}(n)$ is odd for some $p_2 \equiv 5 \pmod{12}$, and thus $a(n)$ vanishes.

Now that we have found the Fourier coefficients of $qE^2(q^{12})$, we turn our attention to discriminant $-576 = -144 \cdot 4$. We have $\mathrm{CL}(-576) \cong C_4 \times C_2$ and we list the reduced forms according to their genus

$\mathrm{CL}(-576) \cong C_4 \times C_2$		$\left(\frac{p}{3}\right)$	$\left(\frac{2}{p}\right)$	$\left(\frac{-1}{p}\right)$
Principal Genus	$(1, 0, 144)$, $(9, 0, 16)$	$+1$	$+1$	$+1$
Second Genus	$(4, 4, 37)$, $(13, 10, 13)$	$+1$	-1	$+1$
Third Genus	$(5, 2, 29)$, $(5, -2, 29)$	-1	-1	$+1$
Fourth Genus	$(9, 6, 17)$, $(9, -6, 17)$	-1	$+1$	$+1$

In the above table, p is taken to be coprime to -576 and represented by the given genus.

We write (5.9) as

$$\frac{\phi(q)\phi(q^{36}) - \phi(q^4)\phi(q^9)}{2} = qE^2(q^{12}). \tag{5.14}$$

Applying (1.13) to the left side of (5.14), we obtain

$$\psi(q^8)\phi(q^{36}) - q^8\phi(q^4)\psi(q^{72}) = E^2(q^{12}). \tag{5.15}$$

Replacing q^4 by $-q^4$ in (5.15), yields

$$\psi(q^8)\phi(-q^{36}) - q^8\phi(-q^4)\psi(q^{72}) = E^2(-q^{12}). \tag{5.16}$$

Multiplying (5.16) by q gives

$$q\psi(q^8)\phi(-q^{36}) - q^9\phi(-q^4)\psi(q^{72}) = qE^2(-q^{12}). \tag{5.17}$$

Taking $d = -576$ and $m = 1, 9$ in Theorem 2.2, we obtain

$$\frac{B(1, 0, 144, q) - B(4, 4, 37, q)}{2} = q\phi(-q^{36})\psi(q^8), \tag{5.18}$$

and

$$\frac{B(9, 0, 16, q) - B(13, 10, 13, q)}{2} = q^9\phi(-q^4)\psi(q^{72}). \tag{5.19}$$

The difference between (5.18) and (5.19) is exactly the left side of (5.17). Hence,

$$\frac{B(1,0,144,q)-B(4,4,37,q)-B(9,0,16,q)+B(13,10,13,q)}{2}=qE^2(-q^{12}). \tag{5.20}$$

One can verify

$$qE^2(-q^{12})=q\frac{E^6(q^{24})}{E^2(q^{12})E^2(q^{48})}, \tag{5.21}$$

which is shown to be an eigenform for all Hecke operators in [13]. Thus we have the relation

$$[q^n]qE^2(-q^{12})=\begin{cases}[q^n]qE^2(q^{12}) & \text{if } n\equiv 1 \pmod{24},\\ -[q^n]qE^2(q^{12}) & \text{if } n\equiv 13 \pmod{24},\\ 0 & \text{if } n\not\equiv 1 \pmod{12}.\end{cases} \tag{5.22}$$

Employing (5.22) along with (5.13) yields a formula for the Fourier coefficients of $qE^2(-q^{12})$.

6. Concluding Remarks

The multiplicative combinations contained in the examples of sections 3, 4, and 5 have the fortunate property of being eigenforms for all Hecke operators. We briefly outline an example that does not share the mentioned property.

Applying Theorem 2.2 to discriminant -1152 yields

$$\frac{B(1,0,288,q)-B(4,4,73,q)}{2}=q\psi(q^8)\phi(-q^{72}), \tag{6.1}$$

and

$$\frac{B(9,0,32,q)-B(17,2,17,q)}{2}=q^9\psi(q^{72})\phi(-q^8), \tag{6.2}$$

and we see the products (6.1) and (6.2) are not disjoint. We are not be able to separate these products by congruences.

We have $\mathrm{CL}(-1152)\cong C_4\times C_2$ and the reduced forms listed by genus are

$\mathrm{CL}(-1152)\cong C_4\times C_2$		$\left(\frac{p}{3}\right)$	$\left(\frac{2}{p}\right)$	$\left(\frac{-1}{p}\right)$
Principal Genus	$(1,0,288)$, $(4,4,73)$	$+1$	$+1$	$+1$
Second Genus	$(9,0,32)$, $(17,2,17)$	-1	$+1$	$+1$
Third Genus	$(11,6,27)$, $(11,-6,27)$	-1	-1	-1
Fourth Genus	$(16,8,19)$, $(16,-8,19)$	$+1$	-1	-1

.

In the above table, p is taken to be coprime to -1152 and represented by the given genus.

Using the fifth row of Table 1 with $A=(11,6,27), B=(9,0,32)$, $\langle A,B\rangle\cong C_4\times C_2$, we have

$$\theta_+(q):=\frac{B(1,0,288,q)-B(4,4,73,q)+B(9,0,32,q)-B(17,2,17,q)}{2}, \tag{6.3}$$

and

$$\theta_-(q):=\frac{B(1,0,288,q)-B(4,4,73,q)-(B(9,0,32,q)-B(17,2,17,q))}{2}, \tag{6.4}$$

are both multiplicative. We find that $\theta_-(q)$ is an eigenform for all Hecke operators, yet $\theta_+(q)$ is an eigenform for all Hecke operators **except** T_3. Unlike previous examples, we are not able to employ (1.18) to find $[q^{3^k}]\theta_+(q)$. However, one can show

$$T_3(\theta_+(q))=\frac{B(1,0,32,q^3)-B(4,4,9,q^3)}{2}. \tag{6.5}$$

By Theorem 2.1, or by Theorem 2.2, we have

$$\frac{B(1,0,32,q^3)-B(4,4,9,q^3)}{2}=q^3E(q^{24})E(q^{48}). \tag{6.6}$$

Using the above remarks we are able to derive $[q^{3^k}]\theta_+(q)$. Proceeding similarly to previous examples, one may derive the Fourier coefficients of (6.1) and (6.2).

This result along with other similar results will be discussed elsewhere.

7. Acknowledgement

Acknowledgement. We are grateful to Hamza Yesilyurt, John Voight, and Li-Chien Shen for helpful discussions, and Keith Grizzell for a careful reading of the manuscript. We would like to thank Jean-Pierre Serre for bringing references [4], [9], and [15] to our attention.

References

[1] A. Berkovich and H. Yesilyurt, *Ramanujan's identities and representation of integers by certain binary and quaternary quadratic forms*, Ramanujan J. **20** (2009), no. 3, 375–408

[2] A. Berkovich and H. Yesilyurt, *On Rogers-Ramanujan functions, binary quadratic forms and eta-quotients*, arXiv:1204.1092 [math.NT], accepted for publication in Proc. of A.M.S.

[3] B. C. Berndt,*Ramanujan's Notebooks, Part III*, Springer, New York, 1991.

[4] F. Van Der Blij, *Binary quadratic forms of discriminant* -23, Neder. Akad. Wetensch, **55** (1952), 498–503.

[5] H. H. Chan, S. Cooper, W-C. Liaw, *On* $\eta^3(a\tau)\eta^3(b\tau)$ *with* $a+b=8$, J. Aust. Math. Soc. **84** (2008), 301–313.

[6] O. X. M. Yao, S. Cooper, E. X. W. Xia, J. Jin, *Explicit Formulas for the Fourier Coefficients of a Class of Eta Quotients*, Int. J. Number Theory, **09** (2013), 487–503.

[7] D. A. Cox,*Primes of the Form* x^2+ny^2*: Fermat Class Field Theory and Complex Multiplication*, John Wiley& Sons, New York, 1989.

[8] L. E. Dickson, *Introduction to the Theory of Numbers*, Dover, New York, 1957.

[9] R. Fricke, *Lehrbuch der Algebra*, vol III, Friedr. Vieweg, 1924.

[10] B. Gordon, K. Hughes, *Multiplicative properties of* η *-products II*, Contemporary Mathematics **143** (1993), 415–430.

[11] E. Hecke, *Mathematische Werke*, Vandenhoeck & Ruprecht, Göttingen, 1970.

[12] G. Köhler, *Eta Products and Theta Series Identities*, Springer, New York, 2011.

[13] Y. Martin, *Multiplicative* η *-quotients*, Trans. Amer. Math. Soc. **348** (1996), no. 12, 4825–4856.

[14] A. Okamoto, *On expressions of theta series by* η*-products.* Tokyo J. Math. **34** (2011), no. 2, 319–326.

[15] B. Schoeneberg, *Bemerkungen über einige klassen von modulfunktionen.* Neder. Akad. W. Proc., A, **70** (1967), 177–182.

[16] J-P. Serre, *Modular forms of weight one and Galois representations.* In: A. Froelich ed. Algebraic Number Fields, (1977), 193–268.

[17] C. L. Siegel, *Über die analytische theory der quadratische Formen.* Ann. of Math, **36** (1935), 527–606.

[18] Z. H. Sun, K. S. Williams, *On the number of representations of* n *by* $ax^2+bxy+cy^2$, Acta Arith. **122** (2006), no. 2, 101–171.

[19] J. Voight, *Quadratic forms that represent almost the same primes*, Math. Comp. **76** (2007), 1589–1617.

[20] K. S. Williams, *Fourier series of a class of eta quotients.*, Int. J. Number Theory, **08** (2012), 993–1004.

Department of Mathematics, University of Florida, 358 Little Hall, Gainesville FL 32611, USA
E-mail address: `alexb@ufl.edu`

Department of Mathematics, University of Florida, 358 Little Hall, Gainesville FL 32611, USA
E-mail address: `frankpatane@ufl.edu`